

## Math 2001: PHW8

1. Consider the following

**Claim.** *The number  $n(n+1)$  is an odd number for every  $n$ .*

*Proof.* Assume the statement is true for  $n$ . We prove the statement for  $n+1$  by induction. Note that

$$(n+1)((n+1)+1) = n(n+1) + 2(n+1).$$

By induction  $n(n+1)$  is odd. Thus,  $(n+1)((n+1)+1)$  is the sum of an odd number  $n(n+1)$  and an even number  $2(n+1)$ . The sum of an odd number and an even number is odd. Thus, we have proved the claim by induction.  $\square$

I checked the claim and it doesn't seem to work for  $n = 15$ . What is wrong with the proof?

2. For each of the following sequences,

- Give a formula for the  $n$ th term in the sequence,
- Give a recursive definition for the sequence (ie. initial values and a recursive equation).

(a)  $1, 2, 3, 4, 5, \dots$

(b)  $1, 2, 4, 8, 16, 32, \dots$

(c)  $1, 2, 6, 24, 120, \dots$

3. Let  $f_0, f_1, \dots$  be the Fibonacci sequence. For each of the following

- Decide whether the identity is easier to prove by induction or directly using Binet's formula (and some algebra). Explain.
- Prove the identity using your preferred method.

(a)  $\sum_{k=1}^n f_k^2 = f_n f_{n+1}$ .

(b)  $\sum_{k=0}^n f_k = f_{n+2} - 1$ .

(c)  $f_{2n+1} = f_{n+1}^2 + f_n^2$ .

4. The Lucas sequence is given by

$$L_1 = 1, \quad L_2 = 3, \quad L_n = L_{n-1} + L_{n-2}, \quad n \geq 3.$$

- (a) Find the first 6 values of the Lucas sequence.
- (b) What should  $L_0$  be defined to be to not mess up the recursion?
- (c) Use induction to prove that

$$L_n = f_{n-1} + f_{n+1}, \quad \text{for } n \geq 1,$$

where  $f_n$  is the  $n$ th Fibonacci number.

- (d) Prove that

$$L_n = \left( \frac{1 + \sqrt{5}}{2} \right)^n + \left( \frac{1 - \sqrt{5}}{2} \right)^n.$$